

Factoring Summary

- (1) factor out the Greatest Common Factor (GCF) *Form for GCF:* $ax^2 + bx + ab = a(x^2 + x + b)$
- (2) factor by grouping (see example below)
- (3) form: $\frac{x + b}{x + c} \cdot \frac{ax}{P}$ Find factors of c that add to get b and multiply to get c. 2
- (4) form: $\frac{x^2 + bx + c}{P^2}$ Use trial and error to find the factored form.
- (5) form: $\pm 2PQ + Q^2$ Then this factors into: $(P \pm Q)^2$ (called "perfect square")
- (6) form:

$x^2 - y^2 = (x + y)(x - y)$

- (7) form: *cannot be factored with integers!*

$x^2 + y^2$	$($
-------------	-----
- (8) form:

$x^3 + y^3 = (x + y)(x^2 - xy + y^2)$

- (9) form:

$x^3 - y^3 = (x - y)(x^2 + xy + y^2)$

Examples: Directions - factor all of the following completely.

- (1) $3x^2 + 9x + 15$ has a GCF of 3. (NOTE: ALWAYS FACTOR OUT GCF FIRST!!)
 Thus, factoring out 3 yields: $3(x^2 + 3x + 5)$
(Since the expression inside the parentheses cannot be factored, this is the final answer.)
- (2) $3x^3 + 2x^2 - 6x - 4$ is a candidate for factoring by grouping. Grouping terms:
 $(3x^3 + 2x^2) + (-6x - 4) = x^2(3x + 2) - 2(3x + 2) = (3x + 2)(x^2 - 2)$
- (3) $x^2 + 4x - 12$ Since a=1 in the trinomial, need to find factors of -12 that add to get 4.
 All the possible pairs of factors for -12 are: 1,-12; -1,12; 2,-6; **-2,6**; 3,-4; -3,4 Since the only pair that adds to 4 is {-2,6} the answer is: **$(x - 2)(x + 6)$**
- (4) $3x^2 + 2x - 8$ Since a≠1 in the trinomial, use trial and error to find the answer.
 The factor pairs of 3 are: {3,1}. The factor pairs for -8 are: {1,-8}, {-1,8}, {2,-4}, {-2,4} By trial and error it is found that the answer is **$(3x - 4)(x + 2)$**

<u>Sign Hints</u> :		
If trinomial has the form:	$ax^2 + bx + c$	then factored form is then $(px + m)(qx + n)$
If trinomial has the form:	$ax^2 - bx + c$	factored form is then $(px - m)(qx - n)$
If trinomial has the form:	$ax^2 \pm bx - c$	factored form is $(px + m)(qx - n)$ OR $(px - m)(qx + n)$

(5) $4x^2 - 12x + 9$ is in the form $P^2 \pm 2PQ + Q^2$. Thus, $4x^2 - 12x + 9 = (2x - 3)^2$ (*perfect square*)

(6) $9x^2 - 36y^2$ is in the form of $x^2 - y^2 = (x+y)(x-y)$. Thus, $9x^2 - 36y^2 = (3x + 6y)(3x - 6y)$

(7) $9x^2 + 36y^2$ is in the form of $x^2 + y^2$. Thus, $9x^2 + 36y^2$ is **non-factorable** using integers.

(8) $8x^3 + 27y^3$ is in the form of $x^3 + y^3 = (x+y)(x^2 - xy + y^2)$. Thus, $8x^3 + 27y^3 = (2x + 3y)(4x^2 - 6xy + 9y^2)$

(9) $8x^3 - 27y^3$ is in the form of $x^3 - y^3 = (x-y)(x^2 + xy + y^2)$. Thus, $8x^3 - 27y^3 = (2x - 3y)(4x^2 + 6xy + 9y^2)$